

## FINAL YEAR UNIVERSITY EXAMINATION 2019-2020 Final Year B.Pharm. Semester VIII SUBJECT-BPH\_C\_801\_T-Pharmaceutical Chemistry III **MULTIPLE CHOICE QUESTIONS: QUESTION BANK**

- 1)Identify selective  $\alpha_1$  antagonist containing quinazoline nucleus from the following agents.
- a) Tolazoline b) Phentolamine c) Phenoxybenzamine d) Prazosin
- 2) Mechanism of action of ramelteon is
- a) NMDA receptor antagonist
- b) Melatonin receptor antagonist
- c) GABA receptor antagonist

- d) Serotonin receptor antagonist
- 3) Which of the following Acetylcholine esterase inhibitor does not contain quaternary ammonium group?
- a) Physostigmine
- b) Neostigmine
- c) Pyridostigmined) Edrophonium
- 4) Amitriptyline (structure given below) undergoes metabolism to give active metabolite by



- a) O-demethylation hydroxylation
- b) N-demethylation
- c) Oxidative deamination d) Oxidative
- 5) Which of the following is not a structural feature of opioid antagonist?
- a) Presence of allyl/cyclopropylmethyl group at 17<sup>th</sup> position
- b) Replacement of 6 –OH with Keto group
- c) Presence of 7-8 double bond
- d) Substitution of 14β-OH
- 6) The enzyme which degrades dopamine is
- a) COMT b) DOPA decarboxylase c) Aldehyde reductase d) Cyclooxygenase
- 7) Identify the Non-Steroidal Anti-inflammatory drug that can be synthesized using following reagent.



- a) Diclofenac
- b) Naproxen
- c) Piroxicam
- d) Nimesulide
- 8) Which one of the following statement of trimethadione is inappropriate?
- a) Trimethadione is a prodrug
- b) Trimethadione is used in absence seizures
- c) Trimethadione belongs to imminostilbene class
- d)Trimethadione is antagonist of T type of calcium channels
- 9) ----is purely anabolic steroid with no androgenic activity.
- a) Testosterone
- b) Stanozolol c) 17α-Methyltestosteroned) Oxymesterone
- 10) An antipsychotic drug fluphenazine is given as an ester of decanoic acid
- a) To increase potency

- b) To increase duration of action
- c) To increase chemical stability
- d) To increase binding modes with the receptor
- 11) Name a NSAID associated with minimal gastric side effects
- a) Diclofenac b)Flurbiprofen
- c) Aspirin
- d)Nabumetone
- 12) Which of the following is not tricyclic antidepressant?
- a) Fluoxetine b) Imipramine c) Chlorimipramine d) Amitriptyline
- 13) Which one of the following drug inhibit xanthine oxidase
- a) Febuxostat b) Probenecid
- c) Colchicine
- d) Sulfinpyrazone
- 14) ---- co-administered with levodopa.
- a) Selegline b) Amantadine
- c) Carbidopa
- d) Benztropine
- 15) Which one of the substituent can be present on nitrogen in narcotic antagonist?
- a) Methyl b) Cyclopropyl methyl c) Phenyl ethyl d) Ethyl
- 16) Select an anticonvulsant from the following which acts by Ca<sup>+2</sup> blockade.
- a) Phenytoin
- b) Carbamazepine
- c) Valproric Acid d) Trimethadione

- 17) Identify metabolite of diazepam from the following.a) Chlordiazepoxideb) Oxazepamc) Nitrazepamd) Clonazepam
- 18) Which drug is COMT resistant?
- a) Isoproternol b) Salbutamol c) Colterol d)Epinephrine
- 19) What is relation between xylometazoline and oxymetazoline?
- a)Both are  $\alpha_2$  adrenergic agonists. b) Both are 2-aminoimidazolines
- c) Oxymetazoline is metabolite of xylometazolined)Both are adrenergic antagonists
- 20) Mention therapeutic use of

- a) Mysthenia Gravis b) Insecticide c) Glaucoma d)Alzheimer disease
- 21) Which of the following statement is not true with respect to Imipramine?
- a) It is MAO inhibitor
- b) It gets metabolized by N-dealkylation
- c) It is norepinephrine and serotonin reuptake inhibitor
- d) Dibenzazepine is basic scaffold present
- 22) Succinylcholine needs to be administered with ----- to prevent enzymatic degradation
- a) Tubocurarine b) Gallamine c) Decamethoniumd) Neostigmine
- 23) Why  $17\alpha$  substituent in androstane is important?
- a) Increase potency b) Increase anabolic activity
- c) Provide oral bioavailability d)Increaseprogestational activity
- 24) Mechanism of action of

a)Nonselective β agonist

b) Nonselective β antagonist

c) Selective β<sub>2</sub> agonist

d) selective β<sub>2</sub> antagonist

25) The IUPAC name of Buspirone is

a) 8-[4-(4-pyrimidin-2-yl piperazin-1-yl)butyl}-8-azaspiro [4.5] decane-7,9-dione

b) 8-[4-(4-pyrimidin-2-yl piperazin-1-yl)propyl}-8-azaspiro [4.5] decane-7,9-dione

c) 8-[4-(4-pyridin-2-yl piperazin-1-yl)butyl}-8-azaspiro [4.5] decane-7,9-dione

d) 8-[4-(4-pyrimidin-2-yl piperazin-1-yl)butyl}-8-azaspiro [4.5] decane-6,8-dione

26) Which one of the following anti epileptic drug do not contain ureide in its structure

a) Phenytoin b) Trimethadione c) Ethotoin d)Lamotrigine

27) Refer the general structural of direct acting sympathomimetic. Converting catechol to resorcinol is responsible for------. Select the correct answer.

a) Increased  $\alpha$  receptor selectivity

b)Nonselective nature of drug

c) Increases β<sub>2</sub> receptor selectivity and bioavailability

d) Indirect activity

28) In the structure of Acetylcholine (shown below), substitution of  $\alpha$ -methyl group-----.


a) Increases muscarinic activity

b) Increases nicotinic activity

c) Increases antagonistic activity

d) No change in activity


29) Identify the encircled nucleus in the following.




- a) Dibenzazepine b) Benzodiazepine c) Phenothiazine d) Benzoxazine
- 30) Which drug is a structural hybrid of meperidine and methadone?
- a) Loperamide b) Diphenoxylate c) Fentanyl d) Pheneridine
- 31) In the case of thioxanthene type of an antipsychotics more active isomers have ------configuration.
- a) Cis b) Trans c) E d) R
- 32) Which one of the following drug shows minimal EPS
- a) Pimozide b) Clozapine c) Droperidol d)Chlorpromazine
- 33) Codeine is-----.
- a) Methylated analog of morphine
- b) Acetylated analog of morphine

c) Antagonist of morphine

- d) Metabolite of morphine
- 34) The  $\beta$  blocker which contain arylethanolamine scaffold
- a) Esmolol b) Metoprolol c) Sotalol d) Timolol
- 35) To which chemical class does the anticholinergic drug dicyclominebelong?
- a) Aminoalcoholb) Aminoamides c)Amino alcohol esters d) Amino alcohol ethers
- 36) The given molecule has how many chiral centers?



- 37) Which substituent in celecoxib is responsible precluding binding with COX I enzyme?
- a) Sulfamoyl b) Phenyl
  - c) Methyl d)Trifluromethyl
- 38) Read the statements given below related to SAR of glucocorticoids and choose the correct option given below.



A: Introduction of 9α-Fluorodecreases salt retention property

B: Methyl group at 16<sup>th</sup> position can be either  $\alpha$  or  $\beta$  for better glucocorticoidal activity

a) A correct

b) B correct

c) A and B both correct

d) A and B both wrong

- 39) Which structural feature of morphine is missing in levorphanol?
- a) Phenyl ring b) Epoxide bridge c) Piperidine ring
- d) 03- OH group
- 40) In case of Phenothiazine -Cl group at 2<sup>nd</sup> position and 3 C chain separating N10 and amino nitrogen is essential because ------.

a) It is important for lipophilicity

b) It gives dopamine like arrangement

c) To maintain potency

d) It increases CNS penetration.

41) Center of acidity in case of Piroxicam can be represented by---

a) -SO<sub>2</sub>NCH<sub>3</sub> group b)Enolic acid c) Pyridine ring d) –OH group

42) Identify the incorrect pair from the following.

a) Fluoxetine: Phenoxyphenylpropylamine

b) Vigabtarine: γ-Vinyl GABA

c) Bethanechol: Carbamate ester

d)Ritodrine: Catechol hydroxyl

- 43) Which one of the following group can be tolerated in 7<sup>th</sup> position of benzodiazepine ring
- a) -NH<sub>2</sub> b)-OH c) -NO<sub>2</sub> d) -CH<sub>3</sub>
- 44) Find the wrong pair
- a) Prazosin :  $\alpha_1$  blocker b) Guanabenz:  $\alpha_2$  agonist
- c) Phenylephrine:  $\alpha_1$  agonist d)Phenylephrine:  $\alpha_2$  antagonist
- 45) Identify the correct statement related to benzomorphan.
- a) They are obtained by removal of epoxide bridge of morphine.
- b) They do not contain B, C, E ring of morphine
- c) Cis and Trans isomer have unequal analgesic activity
- d) They are obtained by removal of C ring of morphine
- 46) Find the odd one out
- a) Gallamine b) Succinylcholine
- c) Decamethonium
- d) Procyclidine
- 47) Which of the following benzodiazepine is suitable for treatment in children & old age patients?
- a) Diazepam b) Oxazepam c) Temazepam d) Nitrazepam
- 48) What is the mechanism of action of

- a) Inhibit reuptake of GABA
- b) Inhibit GABA transaminase
- c) Inhibit glutamate decarboxylase
- d)Blocks GABA receptor
- 49) Modification of nonrigid opioid like ---- leads to butyrophenone whereas butyrophenones are modified to------type of antipsychotic.

Complete the statement by selecting the correct pair from the following.

- a)Meperidine, Diphneylbutylpiperidine
- b) Methadone, Diphneylbutylpiperidine
- c) Meperidine, Benzamide
- d) Methadone: Benzamide
- 50) Which conformer of acetylcholine binds with the receptor?
- a) Synperiplanar
- b) Antiperiplanal
- c) Synclinal
- d) Anticlinal

Q. 51 Which of the following is long acting sedative hypnotic?

A] Diazepam

B] Alprazolam

C] Temazepam

D] Imipramine

Q. 52 Which is the **incorrect** statement with respect to structure given in Q. 4

A] Ring C is ortho substituted with electron withdrawing group for optimum activity

B] Ring C when para substituted increases activity

C] Ring C is diortho substituted with electron withdrawing group for optimum activity

D] Ring C when para substituted decreases activity

Q. 53 Which one of the following is Cytokine inhibitor?

A] Abatacept

B] Fluoxetine

C] Propranolol

D]Aldosterone

Q. 54 Anakinra is

A]costimulation modulator

B] anti-TNF alpha antibody

C] TNF-alpha blocker

D]interleukin receptor antagonist

Q. 55 Alprazolam (structure given below) is short acting sedative hypnotic because of

A] Presence of 4,5 double bond in diazepine ring

B] Presence of 1,4-diazepine ring

C] Presence of chlorine on phenyl ring

D] Presence of methyl group on triazole ring
Q. 56 Nicotine receptor is a type of which receptor?
A] GPCR
B] Ion channel receptor
C] Kinase receptor
D] Nuclear receptor
Q. 57 Donepezil contains which heterocycle as scaffold?
A] Pyridine
B] Piperidine
C]Piperazine
D] Pyrazine
Q. 58 Chemical class of Ibuprofen is,
A] Aryl acetic acid
B] Aryl propanoic acid
C] Indole acetic acid
D]Anthranilic acid
Q. 59 Which one of the following is example of mineralo-corticosteroid
A]betamethasone
B] prednisone
C]Aldosterone
D]triamcinolone
Q. 60 Which of the following is not an example of progestin?
A] Tamoxifen
B]Megestrol
C] Norethindrone
D]Medroxypogesterone
Q. 61 Identify the drug with the following chemical name :
$17\alpha$ -pregna-2,4-dien-20-yno-[2,3- $d$ ]isoxazol-17-ol
a. Stanazolol

- b. Nandrolone
- c. Danazole
- d. Oxandrolone
- Q. 62 Diazepam is metabolized to Nordazepam by \_\_\_\_\_and\_\_\_\_
  - a. CYP2C19 & CYP3A4
  - b. CYP2C18 & CYP3A3
  - c. CYP2C17 & CYP3A2
  - d. CYP2C20 & CYP3A5
- Q. 63 Identify the incorrect statement with reference to the following:

- a. Approved for analgesic therapy and for the maintenance and treatment of opioid addiction.
- b. opioid activity resides in the S-enantiomer
- c.  $\mu$ -receptor agonist
- d. synthetic opioid
- Q. 64  $\alpha$ -CH3 substitutent present in the profens,
  - a. decreases cyclooxygenase inhibitory activity and induces toxicity of the profens.
  - b. decreases cyclooxygenase inhibitory activity and decreases toxicity of the profens.
  - c. increases cyclooxygenase inhibitory activity and increases toxicity of the profens.
  - d. increases cyclooxygenase inhibitory activity and reduces toxicity of the profens.
- Q. 65 What kind of change in the structure of direct acting sympathomimetic are responsible for making them indirectly acting compounds?
- a) Removal of 3'-OH
- b) Removal of 4'-OH
- c) Removal of 3',4'- OH
- d) Retention of 3'-OH and replacing 4'-OH with -CH<sub>2</sub>OH
- Q. 66 Identify the IUPAC nomenclature of following

- a. 7-chloro-1-methyl-5-phenyl-3H-1,4-benzodiazepin-2-one
- b. 7-chloro-2-methylamino-5-phenyl-1,4-benzodiazepine-4-oxide
- c. 7-nitro-5-phenyl-1H-benzo [e] [1,4]diazepin-2(3H)-one
- d. 7-Chloro-1,3-dihydro-1-methyl-5-phenyl-1,4-benzodiazepin-2-one
- Q. 67 Which of the following is aminoalcohol type of an anticholinergic?
- a) Procyclidine b)Benztropine c) Isopropamide d) Tropicamide
- Q. 68 Which is the correct IUPAC name for the following structure?

- a. [(7S)-6-(5-chloropyridin-2-yl)-5-oxo-7H-pyrrolo[3,4-b]pyrazin-7-yl]4-methyl piperazine-1-carboxylate.
- b. [(7S)-6-(5-chloropyridin-2-yl)-5-oxo-7H-pyrrolo[3,4-b]piperazin-7-yl]4-methylpyrazine-1-carboxylate.
- c. [(7S)-6-(5-chloropyridin-2-yl)-5-oxo-7H-pyrrolo[3,4-b]piperidin-7-yl]4-methylpyrimidin-1-carboxylate.
- d. [(7S)-6-(5-chloropyridin-2-yl)-5-oxo-7H-pyrrolo[3,4-b]furan-7-yl]4-methyl piperazine-1-carboxylate.