
ORIENTAL COLLEGE OF PHARMACY

Sem VI, Pharmaceutical Chemistry I (CBCGS) Question Bank (Answers marked in bold) (Total 100 MCQ's), Prepared by: Dr. Nutan Rao


1) What type of molecule is the following structure?

- a) A protein
- b) A nucleic acid
- c) A phospholipid
- d) A carbohydrate
- 2) What is meant by a binding site?
- a) The area of a macromolecular target that is occupied by a drug when it binds.
- b) The portion of the drug to which a drug target binds.
- c) The functional groups used by a drug in binding to a drug target.
- d) The bonds involved in binding a drug to its target.
- 3) Consider the molecule in blue bound to a binding site. Identify the binding interactions taking place at i and iv shown in red.




- a) hydrogen bonds
- b) ionic bonds
- c) van der Waals interactions
- d) dipole-dipole interactions

4) Consider the molecule in blue bound to a binding site. Identify the binding interactions taking place at iii shown in red.



- a) hydrogen bonds
- b) ionic bonds
- c) van der Waals interactions
- 5) Consider the molecule in blue bound to a binding site. Identify the binding interactions taking place at ii shown in red.



- a) hydrogen bonds
- b) ionic bonds
- c) van der Waals interactions
- d) covalent bond
- 6) Which of the following functional groups is most likely to participate in a dipoledipole interaction?
- a) Aromatic ring
- b) Ketone
- c) Alcohol
- d) Alkene

- 7) Which of the following statements is untrue about protein secondary structure?
- a) The alpha helix, beta pleated sheet and beta turns are examples of protein secondary structure.
- b) The ability of peptide bonds to form intramolecular hydrogen bonds is important to secondary structure.
- c) The steric influence of amino acid residues is important to secondary structure.
- d) The hydrophilic/hydrophobic character of amino acid residues is important to secondary structure.

8) Identify which of the following terms refers to the arrangement of different protein subunits in a multiprotein complex.

- a) primary structure
- b) secondary structure
- c) tertiary structure
- d) quaternary structure

9) Which of the following statements is not true about receptors?

- a) Most receptors are proteins situated in the cell membrane.
- b) Receptors contain a hollow or cleft on their surface which is known as a binding site.
- c) Receptors bind chemical messengers such as neurotransmitters or hormones.
- d) Receptors catalyse reactions on chemical messengers.

10) Which of the following is not a neurotransmitter?

- a) acetylcholine
- b) cyclic AMP
- c) noradrenaline
- d) dopamine

11) Which of the following statements is not true about a ligand-gated ion channel receptor?

- a) Ligand-gated ion channel receptors are present in the cell membrane.
- b) Neurotransmitters can act as the chemical messengers for ligand-gated ion channels.
- c) Ligand-gated ion channels consist of five glycoproteins.
- d) Differences in membrane potential affect whether ligand-gated ion channel receptors open or close.

12) Which of the following is not a G-protein coupled receptor?

- a) the muscarinic receptor
- b) the glycine receptor
- c) the adrenergic receptor
- d) the glutamate receptor

13) Which of the following pairs of receptors are likely to show the greatest structural similarity?

- a) the dopamine receptor subtypes D₃ and D₅
- b) the M_2 muscarinic receptor and the β_2 -adrenergic receptor
- c) the H_2 histamine receptor and the α_1 -adrenoceptor
- d) the H_1 histamine receptor and the β_2 adrenoceptor

14) Which of the following reactions is catalysed by a protein kinase?

- a) the phosphorylation of alcohol groups in protein substrates
- b) the hydrolysis of phosphate groups in protein substrates
- c) the phosphorylation of alcohol groups in carbohydrates
- d) the hydrolysis of phosphate groups in ATP and GTP

15) Which of the following is not a typical messenger for a tyrosine kinase linked receptor?

- a) insulin
- b) acetylcholine
- c) growth factors
- d) cytokines

16) Which of the following statements is true regarding the DNA binding region of intracellular receptors?

- a) It contains five cysteine residues.
- b) Four cysteine residues are involved in binding two zinc ions.
- c) It identifies particular nucleotide sequences in DNA.
- d) The DNA binding region is known as having 'thiol fingers'.

17) Which of the following reactions is catalyzed by the enzyme adenylate cyclase?

- a) the conversion of ATP to cyclic AMP
- b) the conversion of cyclic AMP to AMP
- c) the conversion of cyclic AMP to ATP
- d) the conversion of AMP to cyclic AMP

18) What is the product, indicated by the letter C in the following diagram?


a) inositol triphosphate

- b) diacylglycerol
- c) inositol diphosphate
- d) phosphatidylinositol diphosphate

19) Which of the following is true when a G-protein interacts with a receptor?

- a) The G-protein is split into a y-subunit and an α,β -dimer
- b) The G-protein is split into a β -subunit and an α , γ -dimer
- c) The G-protein is split into an α -subunit and a β , γ -dimer
- **d**) The G-protein is split into its component protein subunits

20) What type of structures are the compounds (i) - (iv)?



- a) nucleic acids
- b) nucleotides
- c) nucleosides
- d) deoxyriboses
- 21) To which of the following does guanine form hydrogen bonds in DNA?
- a) adenine
- b) thymine
- c) cytosine
- d) guanine
- 22) Which of the following terms is used to describe a drug that has the same effect on a receptor as the endogenous chemical messenger?
- a) agonist
- b) antagonist
- c) partial agonist
- d) inverse agonist
- 23) Which of the following terms is used to describe a drug that binds to a receptor, and activates it, but to a lesser extent than the endogenous chemical messenger?
- a) agonist
- b) antagonist
- c) partial agonist
- d) inverse agonist
- 24) Which of the following terms is used to describe a drug that binds to a receptor, fails to activate it and leads to a drop in inherent biological activity?
- a) agonist
- b) antagonist
- c) partial agonist
- d) inverse agonist
- 25) Which of the following terms applies to the maximum biological effect resulting from a drug binding to its target?
- a) affinity
- b) efficacy
- c) potency
- d) stability

26) Which of the following terms is the measure of how strongly a drug binds to a receptor?

- a) affinity
- b) efficacy
- c) potency
- d) stability

27) The following antibacterial agent was used in the Second World War.

$$\bigoplus_{H_3N} \bigvee \bigvee_{NH_3} \bigoplus_{NH_3}$$

What is the name of the structure?

- a) Erythromycin
- b) Proflavine
- c) Chloramphenicol
- d) Rifampicin

28) The following structure is a synthetic antibacterial agent called ciprofloxacin.

What is its mechanism of action?

- a) Topoisomerase poison
- b) Metallating agent
- c) Chain terminator
- d) Antisense agent

29) Which of the following statements is the closest description of Phase I metabolism?

- a) Reactions which add a polar molecule to a functional group already present on a drug or one of its metabolites.
- b) Reactions which occur in the blood supply.
- c) Reactions which add a polar functional group to a drug.
- d) Reactions which occur in the gut wall.


30) Which of the following statements is the closest description of Phase II metabolism?

- a) Reactions which add a polar molecule to a functional group already present on a drug or one of its metabolites.
- b) Reactions which occur in the blood supply.
- c) Reactions which add a polar functional group to a drug.
- d) Reactions which occur in the gut wall.

31) Which of the following enzymes is not involved in catalysing a Phase I metabolic reaction?

- a) flavin-containing monooxygenases
- b) monoamine oxidases

- c) glucuronyltransferase
- d) esterases
- 32) Which of the following reactions is not a Phase I metabolic transformation?
- a) reduction of ketones
- b) conjugation to alcohols
- c) oxidation of alkyl groups
- d) ester hydrolysis
- 33) Some drugs containing an ester group are inactive *in vitro*, but are active once the drug has been absorbed *in vivo*. What term is used for such drugs?
- a) predrugs
- b) metabolites
- c) prodrugs
- d) predrugs
- 34) Some peptides and proteins have been used as drugs. Which of the following statements is untrue?
- a) Protein drugs suffer a disadvantage in that they could produce an immune response.
- b) Peptides and proteins generally show poor bioavailability.
- c) Peptide drugs are susceptible to peptidase enzymes.
- d) Peptide drugs are susceptible to metabolic enzymes but not to digestive enzymes.
- 35) What crucial feature of a penicillin is involved in its mechanism of action?
- a) Carboxylic acid
- b) β-lactam ring
- c) Acyl side chain
- d) Thiazolidine ring
- 36) What is the target for clavulanic acid?



- a) The transpeptidase enzyme
- b) L-ala racemase
- c) **B-lactamase**
- d) Penicillin acylase
- 37) Which of the following antibiotics is a macrolide?
- a) Chloramphenicol
- b) Doxycycline
- c) Erythromycin
- d) Streptomycin

38) The following structure is a synthetic antibacterial agent.

$$\begin{array}{c|c} & & & \\ & & \\ & & \\ & & & \\ & & \\ & & & \\ & & & \\ & & \\ & & & \\ & & \\ & & & \\ & & \\ & & & \\ & & \\ & &$$

To which group of compounds does the structure belong?

- a) Aminoacridines
- b) Aminoglycosides
- c) Fluoroquinolones
- d) Tetracyclines

39) What sort of receptor is the nicotinic receptor?

- a) A G-protein coupled receptor
- b) A kinase linked receptor
- c) An intracellular receptor
- d) An ion channel

40) What sort of receptor is the muscarinic receptor?

a) A G-protein coupled receptor

- b) A kinase linked receptor
- c) An intracellular receptor
- d) An ion channel

41) Which of the following is a natural chemical messenger for the adrenergic receptor?

- a) Acetylcholine
- b) Dopamine
- c) Serotonin
- d) Nor-adrenaline

42) Which of the following is a β -lactam antibiotic?

- (a) Penicillin + cephalosporin
- (b) Streptomycin + gentamycin
- (c) Minocyclin + doxycycline
- (d) Chloramphenicol

43) 2,6-Dimethoxy phenyl penicillin is IUPAC of

- (a) Methicillin
- (b) Ampicillin
- (c) Amoxicillin
- (d) Carbencillin

44) Peptidoglacan is made up of _____ amino sugar part.

- (a) N-acetyl glucosamine+N-acetyl muramic acid
- (b) N-acetyl biosamine+N-acetyl muramic acid
- (c) N-acetyl glucosamine+ N-acetyl glucosamine

(d) N-acetyl muramic acid+ N-acetyl muramic acid

45) Which floro quinolone does not contain cyclopropane ring at N-1 position

- (a) Gatifloxacin
- (b) Ciprofloxacin
- (c) Sparfloxacin
- (d) Ofloxacin

46) Which of the following is not present in macrolide?

- (a) A large lactone ring
- (b) A glycosidically linked amino sugar
- (c) A spiroketal group
- (d) A ketone group

47) Which of the following is not a synthetic drug?

- (a) Isoniazide
- (b) Rifampin
- (c) Pyrazinamide
- (d) Ethionamide

48) Which enzyme combination is involved in ergosterol biosynthesis?

A) Lanosterol 14alpha demethylase and Squalene epoxidase

- B) Lanosterol epoxidase and Squalene16alpha demethylase
- C) Lanosterol epoxidase and Squalene14alpha demethylase
- D) Lanosterol 16alpha demethylase and Squalene epoxidase

49) Which of these is not a Polyene?

- A) Terbinafine
- B) Nystatin
- C) Amphotericin B
- D) None

50) What is an Allylamines mode of action?

- A) Inhibit ergosterol synthesis via Lanosterol epoxidase
- B) Inhibit ergosterol synthesis via Squalene 14alpha demethylase
- C) Inhibit ergosterol synthesis via Squalene epoxidase
- D) Inhibit ergosterol synthesis via Lanosterol 14alpha demethylase

51) The drug used for malaria chemoprophylaxis and treatment:

- a) Chloroquine
- b) Quinidine
- c) Quinine
- d) Sulfonamides

52) Which of the following antimicrohials has antipseudomonal action:

- a. Cefpodoxime.
- b. Ceforanide.
- c. Cefotetan.

d. Cefoperazone.

- 53) Which of the following agent has trioxane ring?
- (a) Artemether
- (b) Metronidazole
- (c) Halofantrine
- (d) Prongunil

